# INDEX TO VOLUME 26

## Subject Index


### Abortion

Abortion politics and U.S. population aid: coping with a complex new law, Cohen SA, Special Report, p. 137.

Abortion services in South Africa: available yet not accessible to all, Varkey SJ, Viewpoint, p. 87.

Correcting the record on ARAG's philosophy, Dyer M, Letter, p. 43.

Early medical abortion regimens using different dosages of mifepristone are equally successful, Digest, p. 140.

Ship that makes waves, Update, p. 54.

Women show they can do the math, Update, p. 99.

Work in progress: the expansion of access to abortion services in South Africa following legalization, Althaus FAA, Special Report, p. 84.

### Africa

Abortion services in South Africa: available yet not accessible to all, see **Abortion**.

AIDS and economics in Africa, Update, p. 146.

Changes in family-building patterns in Egypt and Morocco: a comparative analysis, see **Fertility: Planning and Effects**.

Community-based distribution in Tanzania: costs and impacts of alternative strategies to improve worker performance, Janowitz B, Chege J, Thompson A, Rutenberg N and Homan R, p. 158.

Condom use in marital and nonmarital relationships in Zimbabwe, Adetunji J, p. 196.

Couples' report of their contraceptive use: do husbands in Africa overstate the case? see **Males**.

Do service providers in Tanzania unnecessarily restrict clients' access to contraceptive methods? see **Service Providers**.

Exclusively breastfed infants are less likely than those given supplements to contract HIV from their mother, see **Breastfeeding**.

Gender differences in the timing of first intercourse: data from 14 countries, see **Sexual Behavior**.

HIV-infected partner's viral load is the main factor in the risk of transmission in heterosexual couples, see **Sexually Transmitted Diseases**.

Providers, clients okay emergency contraception in Nairobi and Mexico City, see **Emergency Contraception**.

Reproductive decision-making in the context of HIV and AIDS: a qualitative study in Ndola, Zambia, see **Sexually Transmitted Diseases**.

Severity of genital cutting is linked to complications experienced later in life, see **Reproductive Health**.

Sexual risk behaviors among young people in Bamenda, Cameroon, see **Teenagers**.

Single mothers often stay unwed, Update, p. 2.

STDs decline among South African migrant workers after high-risk local women receive treatment, Digest, see **Sexually Transmitted Diseases**.

Trends in the demand for family limitation in developing countries, see **Fertility: Planning and Effects**.

Work in progress: the expansion of access to abortion services in South Africa following legalization, see **Abortion**.

### Asia

Filipino women who use a modern method prefer the pill or tubal sterilization, see **Contraceptive Practice**.

Gender differences in the timing of first intercourse: data from 14 countries, see **Sexual Behavior**.

Home-based neonatal care by village health workers in rural India reduces deaths from bacterial infection, see **Maternal and Child Health**.

Home visits and family-size norms, see **Fertility: Planning and Effects**.

In India, poverty and lack of education are associated with men's physical and sexual abuse of their wives, see **Males**.

Indian men with higher socioeconomic status are more likely to be knowledgeable about reproductive health, Digest, p. 143.

Self-assessment and peer review: improving Indonesian service providers' communication with clients, see **Service Providers**.

STD rates soar in China; three in four new cases are among the unmarried, see **Sexually Transmitted Disease.**

Trends in the demand for family limitation in developing countries, see **Fertility: Planning and Effects**.

Two approaches to managing vaginal discharge lead to overtreatment, missed infections and wasted funds, see **Program Evaluation and Funding**.

Use of a spacing method before sterilization among couples in Kerala, India, see **Contraceptive Practice**.

Women's exposure to mass media is linked to attitudes toward contraception in Pakistan, India and Bangladesh, Digest, p. 48.

### Barrier Methods

At Nicaraguan motels rented for sexual encounters, making condoms available in rooms increases use, Digest, p. 201.

Condom use in marital and nonmarital relationships in Zimbabwe, see **Africa.**

Is the diaphragm a viable option for women in Turkey? Ortayli N, Bulut A, Nalbant H and Cottingham J, p. 36.

### Breastfeeding

Exclusively breastfed infants are less likely than those given supplements to contract HIV from their mother, Digest, p. 46.

Most infant HIV infection from breast milk occurs within six weeks of birth, see **Maternal and Child Health.**

Safe method for nursing women, see **Contraceptive Research**.

Support for longer breastfeeding, Update, p. 2.

### Contraceptive Practice

Contraceptives for the wealthy? Update, p. 99.

Couples' reports of their contraceptive use: do husbands in Africa overstate the case? Digest, p. 203.

Effectiveness of a standard-rule method of calendar rhythm among Mayan couples in Guatemala, see **Natural Family Planning**.

Fertility preferences and contraceptive change in developing countries, **Fertility: Planning and Effects**.

Filipino women who use a modern method prefer the pill or tubal sterilization, Digest, p. 93.

Is the diaphragm a viable option for women in Turkey? see **Barrier Methods**.

Islamic precepts and family planning: the perceptions of Jordanian religious leaders and their constituents, see **Religious Attitudes**.

Use of a spacing method before sterilization among couples in Kerala, India, Zavier F and Padmadas SS, p. 29.

### Contraceptive Research

Once-a-month injectable is acceptable, offers effective protection from pregnancy, see **Hormonal Implants and Injectables**.

Providers, clients okay emergency contraception in Nairobi and Mexico City, see **Emergency Contraception**.

Safe method for nursing women, Update, p. 2.

### Emergency Contraception

Emergency contraception in Ghana, Update, p. 54.

Providers, clients okay emergency contraception in Nairobi and Mexico City, Digest, p. 95.

### Europe

Gender differences in the timing of first intercourse: data from 14 countries, see **Sexual Behavior**.

### Fertility: Planning and Effects

Changes in family-building patterns in Egypt and Morocco: a comparative analysis, Eltigani EE, p. 73.

Effects of an entertainment-education radio soap opera on family planning and HIV prevention in St. Lucia, see **Program Evaluation and Funding**.

# INDEX TO VOLUME 26

## Subject Index


### Abortion

Abortion politics and U.S. population aid: coping with a complex new law, Cohen SA, Special Report, p. 137.

Abortion services in South Africa: available yet not accessible to all, Varkey SJ, Viewpoint, p. 87.

Correcting the record on ARAG's philosophy, Dyer M, Letter, p. 43.

Early medical abortion regimens using different dosages of mifepristone are equally successful, Digest, p. 140.

Ship that makes waves, Update, p. 54.

Women show they can do the math, Update, p. 99.

Work in progress: the expansion of access to abortion services in South Africa following legalization, Althaus FAA, Special Report, p. 84.

### Africa

Abortion services in South Africa: available yet not accessible to all, see **Abortion**.

AIDS and economics in Africa, Update, p. 146.

Changes in family-building patterns in Egypt and Morocco: a comparative analysis, see **Fertility: Planning and Effects**.

Community-based distribution in Tanzania: costs and impacts of alternative strategies to improve worker performance, Janowitz B, Chege J, Thompson A, Rutenberg N and Homan R, p. 158.

Condom use in marital and nonmarital relationships in Zimbabwe, Adetunji J, p. 196.

Couples' report of their contraceptive use: do husbands in Africa overstate the case? see **Males**.

Do service providers in Tanzania unnecessarily restrict clients' access to contraceptive methods? see **Service Providers**.

Exclusively breastfed infants are less likely than those given supplements to contract HIV from their mother, see **Breastfeeding**.

Gender differences in the timing of first intercourse: data from 14 countries, see **Sexual Behavior**.

HIV-infected partner's viral load is the main factor in the risk of transmission in heterosexual couples, see **Sexually Transmitted Diseases**.

Providers, clients okay emergency contraception in Nairobi and Mexico City, see **Emergency Contraception**.

Reproductive decision-making in the context of HIV and AIDS: a qualitative study in Ndola, Zambia, see **Sexually Transmitted Diseases**.

Severity of genital cutting is linked to complications experienced later in life, see **Reproductive Health**.

Sexual risk behaviors among young people in Bamenda, Cameroon, see **Teenagers**.

Single mothers often stay unwed, Update, p. 2.

STDs decline among South African migrant workers after high-risk local women receive treatment, Digest, see **Sexually Transmitted Diseases**.

Trends in the demand for family limitation in developing countries, see **Fertility: Planning and Effects**.

Work in progress: the expansion of access to abortion services in South Africa following legalization, see **Abortion**.

### Asia

Filipino women who use a modern method prefer the pill or tubal sterilization, see **Contraceptive Practice**.

Gender differences in the timing of first intercourse: data from 14 countries, see **Sexual Behavior**.

Home-based neonatal care by village health workers in rural India reduces deaths from bacterial infection, see **Maternal and Child Health**.

Home visits and family-size norms, see **Fertility: Planning and Effects**.

In India, poverty and lack of education are associated with men's physical and sexual abuse of their wives, see **Males**.

Indian men with higher socioeconomic status are more likely to be knowledgeable about reproductive health, Digest, p. 143.

Self-assessment and peer review: improving Indonesian service providers' communication with clients, see **Service Providers**.

STD rates soar in China; three in four new cases are among the unmarried, see **Sexually Transmitted Disease.**

Trends in the demand for family limitation in developing countries, see **Fertility: Planning and Effects**.

Two approaches to managing vaginal discharge lead to overtreatment, missed infections and wasted funds, see **Program Evaluation and Funding**.

Use of a spacing method before sterilization among couples in Kerala, India, see **Contraceptive Practice**.

Women's exposure to mass media is linked to attitudes toward contraception in Pakistan, India and Bangladesh, Digest, p. 48.

### Barrier Methods

At Nicaraguan motels rented for sexual encounters, making condoms available in rooms increases use, Digest, p. 201.

Condom use in marital and nonmarital relationships in Zimbabwe, see **Africa.**

Is the diaphragm a viable option for women in Turkey? Ortayli N, Bulut A, Nalbant H and Cottingham J, p. 36.

### Breastfeeding

Exclusively breastfed infants are less likely than those given supplements to contract HIV from their mother, Digest, p. 46.

Most infant HIV infection from breast milk occurs within six weeks of birth, see **Maternal and Child Health.**

Safe method for nursing women, see **Contraceptive Research**.

Support for longer breastfeeding, Update, p. 2.

### Contraceptive Practice

Contraceptives for the wealthy? Update, p. 99.

Couples' reports of their contraceptive use: do husbands in Africa overstate the case? Digest, p. 203.

Effectiveness of a standard-rule method of calendar rhythm among Mayan couples in Guatemala, see **Natural Family Planning**.

Fertility preferences and contraceptive change in developing countries, **Fertility: Planning and Effects**.

Filipino women who use a modern method prefer the pill or tubal sterilization, Digest, p. 93.

Is the diaphragm a viable option for women in Turkey? see **Barrier Methods**.

Islamic precepts and family planning: the perceptions of Jordanian religious leaders and their constituents, see **Religious Attitudes**.

Use of a spacing method before sterilization among couples in Kerala, India, Zavier F and Padmadas SS, p. 29.

### Contraceptive Research

Once-a-month injectable is acceptable, offers effective protection from pregnancy, see **Hormonal Implants and Injectables**.

Providers, clients okay emergency contraception in Nairobi and Mexico City, see **Emergency Contraception**.

Safe method for nursing women, Update, p. 2.

### Emergency Contraception

Emergency contraception in Ghana, Update, p. 54.

Providers, clients okay emergency contraception in Nairobi and Mexico City, Digest, p. 95.

### Europe

Gender differences in the timing of first intercourse: data from 14 countries, see **Sexual Behavior**.

### Fertility: Planning and Effects

Changes in family-building patterns in Egypt and Morocco: a comparative analysis, Eltigani EE, p. 73.

Effects of an entertainment-education radio soap opera on family planning and HIV prevention in St. Lucia, see **Program Evaluation and Funding**.

Fertility preferences and contraceptive change in developing countries, Feyisetan B and Casterline JB, p. 100.

Home visits and family-size norms, Update, p. 3.

Reproductive decision-making in the context of HIV and AIDS: a qualitative study in Ndola, Zambia, see **Sexually Transmitted Diseases.**

Trends in the demand for family limitation in developing countries, Westoff CF and Bankole A, p. 56.

### Hormonal Implants and Injectables

Changes in bone density from hormonal methods are small and temporary, Digest, p. 202.

Implant is effective for seven years, Update, p. 146.

Once-a-month injectable is acceptable, offers effective protection from pregnancy, Digest, p. 91.

### Intrauterine Devices

Egyptian women who use an IUD have a higher risk of anemia than those who rely on other methods, Digest, p. 142.

Screening for STDs in IUD users, Update, p. 147.

### Latin America and the Caribbean

At Nicaraguan motels rented for sexual encounters, making condoms available for rooms increase use, see **Barrier Methods**.

Consistency of self-reports of sexual activity among young adolescents in Jamaica, see **Teenagers**.

Gender differences in the timing of first intercourse: data from 14 countries, see **Sexual Behavior**.

Effectiveness of a standard-rule method of calendar rhythm among Mayan couples in Guatemala, see **Natural Family Planning**.

Effects of an entertainment-education radio soap opera on family planning and HIV prevention in St. Lucia, see **Program Evaluation and Funding**.

High cesarean section rates in Brazil result in large part from nonclinical factors, see **Pregnancy and Childbirth**.

Measuring family planning service quality through client satisfaction exit interviews, see **Service Providers**.

Providers, clients okay emergency contraception in Nairobi and Mexico City, see **Emergency Contraception**.

Trends in the demand for family limitation in developing countries, see **Fertility: Planning and Effects**.

### Males

Condom use in marital and nonmarital relationships in Zimbabwe, see **Africa**.

Couples' report of their contraceptive use: do husbands in Africa overstate the case? see **Males.**

In India, poverty and lack of education are associated with men's physical and sexual abuse of their wives, Digest, p. 44.

Indian men with higher socioeconomic status are more likely to be knowledgeable about reproductive health, see **Asia**.

Islamic precepts and family planning: the perceptions of Jordanian religious leaders and their constituents, see **Religious Attitudes**.

Unequal time at Kenyan clinics, Update, p. 147.

### Maternal and Child Health

Babies thrive on variety, Update, p. 98.

Exclusively breastfed infants are less likely than those given supplements to contract HIV from their mother, see **Breastfeeding**.

Home-based neonatal care by village health workers in rural India reduces deaths from bacterial infection, Digest, p. 92.

Like mother, like baby, Update, p. 99.

Most infant HIV infection from breast milk occurs within six weeks of birth, Digest, p. 141.

Multiple gestations are associated with adverse outcomes for the mother, Digest, p. 205.

Safe motherhood in Vietnam, Update, p. 54.

Use of antiretroviral treatment pays, Update, p. 98.

### Middle East

Cairo students on FGM, Update, p. 3.

Egyptian women who use an IUD have a higher risk of anemia than those who rely on other methods, see **Intrauterine Devices**.

Is the diaphragm a viable option for women in Turkey? see **Barrier Methods**.

Islamic precepts and family planning: the perceptions of Jordanian religious leaders and their constituents, see **Religious Attitudes**.

Trends in the demand for family limitation in developing countries, see **Fertility: Planning and Effects**.

### Natural Family Planning

Effectiveness of a standard-rule method of calendar rhythm among Mayan couples in Guatemala, Burkhart MC, de Mazariegos L, Salazar S and Lamprecht VM, p. 131.

### Oral Contraceptives

Filipino women who use a modern method prefer the pill or tubal sterilization, see **Contraceptive Practice**.

### Population and Policy Research

City living, Update, p. 98.

### Pregnancy and Childbirth

For fertility, add iodine, Update, p. 98.

High cesarean section rates in Brazil result in large part from nonclinical factors, Digest, p. 47.

Pregnancy, HIV and anemia, Update, p. 55.

### Program Evaluation and Funding

Abortion politics and U.S. population aid: coping with a complex new law, see **Abortion**.

Community-based distribution in Tanzania: costs and impacts of alternative strategies to improve worker performance, see **Africa.**

Effects of an entertainment-education radio soap opera on family planning and HIV prevention in St. Lucia, Vaughan PW, Regis A and St. Catherine E, p. 148.

Implementing the gag rule, Update, p. 2.

Measuring family planning service quality through client satisfaction exit interviews, see **Service Providers**.

Self-assessment and peer review: improving Indonesian service providers' communication with clients, see **Service Providers**.

Two approaches to managing vaginal discharge lead to overtreatment, missed infections and wasted funds, Digest, p. 89.

U.S. law to provide AIDS relief, Update, p. 147.

### Religious Attitudes

Islamic precepts and family planning: the perceptions of Jordanian religious leaders and their constituents, Underwood C, p. 110.

Religion and HIV prevention, Update, p. 146.

### Reproductive Health

Cairo students on FGM, see **Middle East**.

Causes of refugee mortality, Update, p. 3.

Contraceptive need among Cambodian refugees in Khao Phlu camp, Morrison V, p. 188.

Ensuring the reproductive rights of refugees and internally displaced persons: legal and policy issues, Girard F and Waldman W, p. 167.

Indian men with higher socioeconomic status are more likely to be knowledgeable about reproductive health, see **Asia**.

Programmatic responses to refugees' reproductive health needs, Krause SK, Jones RK and Purdin SJ, p. 181.

Reproductive health of war–affected populations: what do we know? McGinn T, p. 174.

Severity of genital cutting is linked to complications experienced later in life, Digest, p. 45.

Turning point: the refugee reproductive health field, Schreck L, Special Report, p. 162.

Why refugees need reproductive health services, Burns K, Malé S and Pierotti D, p. 161.

### Service Providers

Do service providers in Tanzania unnecessarily restrict clients' access to contraceptive methods? Speizer IS, Hotchkiss DR, Magnani RJ, Hubbard B and Nelson K, p. 13.

High cesarean section rates in Brazil result in large part from nonclinical factors, see **Pregnancy and Childbirth**.

Measuring family planning service quality through client satisfaction exit interviews, Williams T, Schutt-Ainé J and Cuca Y, p. 63.

Providers, clients okay emergency contraception in Nairobi and Mexico City, see **Emergency Contraception**.

Self-assessment and peer review: improving Indonesian service providers' communication

with clients, Kim YM, Putjuk F, Basuki E and Kols A, p. 4.

**Sexual Behavior**
Consistency of self-reports of sexual activity among young adolescents in Jamaica, see **Teenagers**.

Gender differences in the timing of first intercourse: data from 14 countries, Singh S, Wulf D, Samara R and Cuca YP, p. 21.

Oral skills needed by sex workers, Update, p. 3.

Sexual risk behaviors among young people in Bamenda, Cameroon, see **Teenagers**.

**Sexually Transmitted Diseases**
AIDS epidemic: an update, Update, p. 3.

HIV-infected partner's viral load is the main factor in the risk of transmission in heterosexual couples, Digest, p. 203.

No to nonoxynol-9 to prevent HIV, Update, p. 147.

Reproductive decision-making in the context of HIV and AIDS: a qualitative study in Ndola, Zambia, Rutenberg N, Biddlecom AE and Kaona FAD, p. 124.

STD rates soar in China; three in four new cases are among the unmarried, Digest, p. 141.

STDs decline among South African migrant workers after high-risk local women receive treatment, Digest, p. 90.

**Status of Women**
Identifying domestic violence, Update, p. 55.

Marriage vs. school in Bangladesh, Update, p. 146.

Women's exposure to mass media is linked to attitudes toward contraception in Pakistan, India and Bangladesh, see **Asia**.

**Sterilization**
Filipino women who use a modern method prefer the pill or tubal sterilization, see **Contraceptive Practice**.

Sterilization as a social strategy? Update, p. 54.

Use of a spacing method before sterilization among couples in Kerala, India, see **Contraceptive Practice**.

**Teenagers**
Consistency of self-reports of sexual activity among young adolescents in Jamaica, Eggleston E, Leitch J and Jackson J, p. 79.

Gender differences in the timing of first intercourse: data from 14 countries, see **Sexual Behavior**.

Sexual risk behaviors among young people in Bamenda, Cameroon, Rwenge M, p. 118.

## Author Index

Adetunji J, Condom use in marital and nonmarital relationships in Zimbabwe, p. 196.

Althaus FA, Work in progress: the expansion of access to abortion services in South Africa following legalization, Special Report, p. 84.

Bankole A, see Westoff CF.

Basuki E, see Kim YM.

Biddlecom, AE, see Rutenberg N.

Bulut A, see Ortayli N.

Burkhart MC, de Mazariegos L, Salazar S and Lamprecht VM, Effectiveness of a standard-rule method of calendar rhythm among Mayan couples in Guatemala, p. 131.

Burns K, Malé S and Pierotti D, Why refugees need reproductive health services, p. 161.

Casterline JB, see Feyisetan B.

Chege J, see Janowitz B.

Cohen SA, Abortion politics and U.S. population aid: coping with a complex new law, Special Report, p. 137.

Cottingham J, see Ortayli N.

Cuca YP, see Singh S; see Williams T.

de Mazariegos L, see Burkhart MC.

Dyer M, Correcting the record on ARAG's philosophy, Letter, p. 43.

Eggleston E, Leitch J and Jackson J, Consistency of self-reports of sexual activity among young adolescents in Jamaica, p. 79.

Eltigani EE, Changes in family-building patterns in Egypt and Morocco: a comparative analysis, p. 73.

Feyisetan B and Casterline JB, Fertility preferences and contraceptive change in developing countries, p. 100.

Girard F and Waldman W, Ensuring the reproductive rights of refugees and internally displaced persons: legal and policy issues, p. 167.

Homan R, see Janowitz B.

Hotchkiss DR, see Speizer IS.

Hubbard B, see Speizer IS.

Jackson J, see Eggleston E.

Janowitz B, Chege J, Thompson A, Rutenberg N and Homan R, Community-based distribution in Tanzania: costs and impacts of alternative strategies to improve worker performance, p. 158.

Jones RK, see Krause SK.

Kaona FAD, see Rutenberg N.

Kim YM, Putjuk F, Basuki E and Kols A, Self-assessment and peer review: improving Indonesian service providers' communication with clients, p. 4.

Kols A, see Kim YM.

Krause SK, Jones RK and Purdin SJ, Programmatic responses to refugees' reproductive health needs, p. 181.

Lamprecht VM, see Burkhart MC.

Leitch J, see Eggleston E.

Magnani RJ, see Speizer IS.

Malé S, see Burns K.

McGinn T, Reproductive health of war-affected populations: what do we know? p. 174.

Morrison V, Contraceptive need among Cambodian refugees in Khao Phlu camp, p. 188.

Nalbant H, see Ortayli N.

Nelson K, see Speizer IS.

Ortayli N, Bulut A, Nalbant H and Cottingham J, Is the diaphragm a viable option for women in Turkey? p. 36.

Padmadas SS, see Zavier F.

Pierotti D, see Burns K.

Purdin SJ, see Krause SK.

Putjuk F, see Kim YM.

Regis A, see Vaughan PW.

Rutenberg N, Biddlecom AE and Kaona FAD, Reproductive decision-making in the context of HIV and AIDS: a qualitative study in Ndola, Zambia, p. 124; see also Janowitz B.

Rwenge M, Sexual risk behaviors among young people in Bamenda, Cameroon, p. 118.

Salazar S, see Burkhart MC.

Samara R, see Singh S.

Schreck L, Turning point: the refugee reproductive health field, p. 162.

Schutt-Ainé J, see Williams T.

Singh S, Wulf D, Samara R and Cuca YP, Gender differences in the timing of first intercourse: data from 14 countries, p. 21.

Speizer IS, Hotchkiss DR, Magnani RJ, Hubbard B and Nelson K, Do service providers in Tanzania unnecessarily restrict clients' access to contraceptive methods? p. 13.

St. Catherine E, see Vaughan PW.

Thompson A, see Janowitz B.

Underwood C, Islamic precepts and family planning: the perceptions of Jordanian religious leaders and their constituents, p. 110.

Varkey SJ, Abortion services in South Africa: available yet not accessible to all, Viewpoint, p. 87.

Vaughan PW, Regis A and St. Catherine E, Effects of an entertainment-education radio soap opera on family planning and HIV prevention in St. Lucia, p. 148.

Waldman W, see Girard F.

Westoff CF and Bankole A, Trends in the demand for family limitation in developing countries, p. 56.

Williams T, Schutt-Ainé J and Cuca Y, Measuring family planning service quality through client satisfaction exit interviews, p. 63.

Wulf D, see Singh S.

Zavier F and Padmadas SS, Use of a spacing method before sterilization among couples in Kerala, India, p. 29.